T5530.CIP (UT-0006)

Inventors:

Rao et al.

Serial No.: Filing Date:

09/109,858 July 2, 1998

Page 2

A Petition for a two (2) month extension of time and the requisite fee are provided herewith. Please enter the following amendments and remarks into the record.

In the Claims:

Please cancel claim 23.

Please amend the claims as follows:

- 12. (amended) A method of isolating a pure population of rodent or human CNS neuron-restricted precursor cells comprising the steps of:
- (a) isolating a population of rodent or human multipotent CNS stem cells which generate both neurons and glia;
- (b) incubating the multipotent CNS stem cells in NEP
 medium;
- (c) replating the multipotent CNS stem cells on laminin in NEP medium in the absence of chick embryo extract to induce cell differentiation;
- (d) removing A2B5+ cells from the differentiating cells via specific antibody capture with an antibody that specifically recognizes A2B5;
- (e) purifying from the supernatant following step (d) a subpopulation of cells expressing embryonic neural cell adhesion

25

T5530.CIP (UT-0006)

Inventors:

Rao et al.

Serial No.: Filing Date:

09/109,858

Page 3

July 2, 1998

molecules via a procedure selected from the group consisting of specific antibody capture, fluorescence activated cell sorting, and magnetic bead capture, wherein said procedure uses an embryonic neural cell adhesion molecule antibody that specifically recognizes polysialated neural cell adhesion molecule (NCAM); and

9 XX

(f) incubating the purified subpopulation of cells in a FGF-containing medium configured for supporting adherent growth thereof to obtain an isolated purified population of rodent or human CNS neuron-restricted precursor cells, wherein said neuron-restricted precursor cells differentiate into CNS neuronal cells upon replacement of adherent growth supporting medium with retinoic acid containing medium and fail to proliferate or differentiate in astrocyte-promoting medium containing FGF and 10% fetal calf serum.

887

21. (amended) A method of isolating a pure population of rodent or human CNS neuron-restricted precursor cells comprising the steps of:

62

- (a) removing a sample of spinal cord tissue from a rodent or human embryo at a stage of embryonic development after closure of the neural tube;
 - (b) dissociating cells comprising the sample of spinal

T5530.CIP (UT-0006)

Inventors:

Rao et al.

Serial No.: Filing Date:

09/109,858 July 2, 1998

Page 4

cord tissue removed from the embryo;

- (c) removing A2B5+ cells from the dissociated cells via specific antibody dapture with an antibody that specifically recognizes A2B5;
- (d) purifying from the supernatant following step (c) a subpopulation expressing embryonic neural cell adhesion molecule via a procedure selected from the group consisting of specific antibody capture, fluorescence activated cell sorting, and magnetic bead capture, using an embryonic neural cell adhesion molecule antibody that specifically recognizes polysialated neural cell adhesion molecule.
- (e) plating the purified subpopulation of cells in feeder-cell-independent culture on a substratum and in a medium configured for supporting adherent growth of the neuron-restricted precursor cells; and
- (f) incubating the plated cells at a temperature and in an atmosphere conducive to growth to obtain an isolated, pure population of neuron-restricted precursor cells, wherein said neuron-restricted precursor cells require FGF for adherent growth, differentiate into CNS neuronal cells upon replacement of adherent growth supporting medium with retinoic acid containing medium and fail to proliferate or differentiate in astrocyte-

Continue of the continue of th

T5530.CIP (UT-0006)

Inventors:

Rao et al.

promoting medium containing FGF and 10% fetal calf serum.

Serial No.: Filing Date:

09/109,858 July 2, 1998

Page 5

(amended) The method of claim 21 wherein said procedure 24. is specific antibody capture.

28. (amended) A method of producing postmitotic neurons from a pure population of neuron-restricted precursor cells comprising:

- (a) culturing a pure population of neuron-restricted precursor cells which require FGF and differentiate into CNS neuronal cells but not into CNS glial cells in proliferating conditions; and
- (b) changing the culture conditions of the neuronrestricted precursor cells from proliferating conditions to differentiating conditions, thereby causing the neuron-restricted precursor cells to differentiate into postmitotic neurons.
- (amended) A method\of isolating a pure population of mouse or human CNS neuron-restricted precursor cells comprising the steps of:
- (a) providing a sample of mouse or human embryonic stem cells:
- (b) removing A2B5+ cells from the sample of mouse or human embryonic stem cells via specific antibody capture with an antibody that specifically recognizes A2B5;

T5530.CIP (UT-0006)

Inventors:

Rao et al.

Serial No.: Filing Date:

09/109,858

Page 6

July 2, 1998

(c) purifying from the supernatant from step (b) a subpopulation expressing embryonic neural cell adhesion molecule via a procedure selected from the group consisting of specific antibody capture, fluorescence activated cell sorting, and magnetic bead capture, using an embryonic cell adhesion molecule antibody specifically recognizes polysialated neural cell adhesion molecule;

(d) plating the purified subpopulation of cells in feeder-cell-independent culture on a substratum and in a medium configured for supporting adherent growth of the neuron-restricted precursor cells; and

(e) incubating the plated cells at a temperature and in an atmosphere conducive to growth of the neuron-restricted precursor cells, wherein said neuron-restricted precursor cells require FGF and differentiate into CNS neuronal cells but not into CNS glial cells.

REMARKS

Claims 12, 15, 16, 21, 23, 24, 26-33 and 59 are pending in the instant application. Claims 12, 15, 16, 21, 23, 24, 26-33 and 59 have been rejected. Claims 12, 21, 24 and 59 have been amended. Claim 23 has been canceled. No new matter has been

(Oy)

and